arXiv:0906.3946v1 [math.CO] 22 Jun 2009

The rainbow k-connectivity of two classes of graphs*

Xueliang Li, Yuefang Sun

Center for Combinatorics and LPMC-TJKLC

Nankai University, Tianjin 300071, P.R. China

E-mail: lxl@nankai.edu.cn; syf@cfc.nankai.edu.cn

Abstract

A path in an edge-colored graph G, where adjacent edges may be colored the same, is called a rainbow path if no two edges of G are colored the same. For a κ-connected graph G and an integer k with $1 \leq k \leq \kappa$, the rainbow k-connectivity $rc_k(G)$ of G is defined as the minimum integer j for which there exists a j-edge-coloring of G such that every two distinct vertices of G are connected by k internally disjoint rainbow paths. Let G be a complete $(\ell+1)$-partite graph with ℓ parts of size r and one part of size p where $0 \leq p < r$ (in the case $p = 0$, G is a complete ℓ-partite graph with each part of size r). This paper is to investigate the rainbow k-connectivity of G. We show that for every pair of integers $k \geq 2$ and $r \geq 1$, there is an integer $f(k,r)$ such that if $\ell \geq f(k,r)$, then $rc_k(G) = 2$. As a consequence, we improve the upper bound of $f(k)$ from $(k+1)^2$ to $ck^{\frac{3}{2}} + C$, where $0 < c < 1$, $C = o(k^{\frac{3}{2}})$, and $f(k)$ is the integer such that if $n \geq f(k)$ then $rc_k(K_n) = 2$.

Keywords: edge-colored graph, rainbow path, rainbow k-connectivity, complete (multipartite) graph

AMS Subject Classification 2000: 05C15, 05C40

1 Introduction

All graphs considered in this paper are simple, finite and undirected. Let G be a nontrivial connected graph with an edge coloring $c : E(G) \to \{1, 2, \cdots, k\}$, $k \in \mathbb{N}$, where

---

*Supported by NSFC, PCSIRT and the "973" program.

adjacent edges may be colored the same. A path of G is called *rainbow* if no two edges of it are colored the same. An edge colored graph G is said to be *rainbow connected* if for any two vertices of G there is a rainbow path of G connecting them. Clearly, if a graph is rainbow connected, it must be connected. Conversely, any connected graph has a trivial edge coloring that makes it rainbow connected, i.e., the coloring such that each edge has a distinct color. Thus, Chartrand et al. [2] defined the *rainbow connection number* of a connected graph G, denoted by $rc(G)$, as the smallest number of colors by using which there is an edge coloring of G such that G is rainbow connected.

Suppose that G is a κ-connected graph with $\kappa \geq 1$. It then follows from a well-known theorem of Whitney that for every integer k with $1 \leq k \leq \kappa$ and every two distinct vertices u and v of G, the graph G contains k internally disjoint $u - v$ paths. Chartrand et al. [2] defined the *rainbow k-connectivity* $rc_k(G)$ of G to be the minimum integer j for which there exists a j-edge-coloring of G such that for every two distinct vertices u and v of G, there exist at least k internally disjoint $u - v$ rainbow paths. Thus by the definition, $rc_1(G) = rc(G)$ is the rainbow connection number of G. By coloring the edges of G with distinct colors, we see that every two vertices of G are connected by κ internally disjoint rainbow paths and that $rc_k(G)$ is defined for every integer k with $1 \leq k \leq \kappa$. Furthermore, $rc_k(G) \leq rc_j(G)$ for $1 \leq k \leq j \leq \kappa$.

The concept of rainbow k-connectivity has applications in communication networks. For details we refer to [2] and [3].

In [2], Chartrand et al. studied the rainbow k-connectivity of the complete graph K_n for various pairs k, n of integers. It was shown in [2] that for every integer $k \geq 2$, there exists an integer $f(k)$ such that $rc_k(K_n) = 2$ for every integer $n \geq f(k)$. They also investigated the rainbow k-connectivity of r-regular complete bipartite graphs for some pairs k, r of integers with $2 \leq k \leq r$, and they showed that for every integer $k \geq 2$, there exists an integer r such that $rc_k(K_{r,r}) = 3$.

This paper is to continue their investigation. We improve the upper bound of $f(k)$ from $(k+1)^2$ to $ck^{\frac{3}{2}} + C$ (here $0 < c < 1$ and $C = o(k^{\frac{3}{2}})$), i.e., from $O(k^2)$ to $O(k^{\frac{3}{2}})$, a considerable improvement. For notations and terminology not defined here, we refer to [1].

2 Main Results

In [2] the following result was obtained.

Theorem 2.1 *([2]) For every integer $k \geq 2$, there exists an integer $f(k)$ such that if $n \geq f(k)$, then $rc_k(K_n) = 2$.* ■

The authors in [2] then obtained an upper bound $(k+1)^2$ for $f(k)$, namely $f(k) \leq (k+1)^2$. In the following we will give a better upper bound for $f(k)$.

Let $K_{\ell[r]}$ denote the complete ℓ-partite graph each part of which contains r elements. Let $V = \bigcup_{s=1}^{\ell} V_s$ where $V_s = \{u_{s,1}, u_{s,2}, \cdots, u_{s,r}\}$ $(1 \leq s \leq \ell)$ is the vertex set of each part, and $U_j = \{u_{1,j}, u_{2,j}, \cdots, u_{l,j}\}$ $(1 \leq j \leq r)$.

We consider the graph $G = K_{\ell^2[r]}$ where ℓ is an integer, and give G a 2-edge-coloring as follows: Let $\overline{V_i} = \bigcup_{t=1}^{\ell} V_{(i-1)\ell+t}$ where $1 \leq i \leq \ell$. We know that the subgraph, denoted by G_j, of G induced by U_j is the complete graph K_{ℓ^2}. Let $U_j = U_{1,j} \cup \cdots \cup U_{\ell,j}$ where $U_{i,j} = \{u_{(i-1)\ell+1,j}, \cdots, u_{i\ell,j}\}$. Then the subgraph, denoted by $G_{i,j}$, of G induced by $U_{i,j}$ is the complete graph K_ℓ. Similar to the edge coloring in the proof of Theorem 2.1 in [2], we first give a 2-edge-coloring of each G_j $(1 \leq j \leq r)$ as follows: We assign the edge uv of G the color 1 if either $uv \in E(G_{i,j})$ for some $i (1 \leq i \leq \ell)$ or if $uv = u_{(i_1-1)\ell+t,j}\ u_{(i_2-1)\ell+t,j}$ for some i_1, i_2, t with $1 \leq i_1, i_2, t \leq \ell$ and $i_1 \neq i_2$. All other edges of G_j are assigned the color 2. With a similar argument to Theorem 2.1 in [2], there are $\ell - 1$ disjoint rainbow $u - v$ paths for any two vertices u, v in the subgraph G_j including 1 path of length 1 and $\ell - 2$ paths of length 2.

For other edges, that is, the edges between distinct G_js, we use above two colors as follows: Let $uv = u_{i_1,j_1}u_{i_2,j_2}$ where $u_{i_1,j_1} \in G_{j_1}, u_{i_2,j_2} \in G_{j_2}$, we give it the color different from the color of edge $u_{i_1,j_1}u_{i_2,j_1}$ of graph G_{j_1} or $u_{i_1,j_2}u_{i_2,j_2}$ of graph G_{j_2} (since edges $u_{i_1,j_1}u_{i_2,j_1}$ and $u_{i_1,j_2}u_{i_2,j_2}$ have the same color by above coloring).

Next we will count the number of disjoint rainbow $u - v$ paths for any two vertices u, v of G with above edge coloring. Without loss of generality, let $u = u_{1,1}$. We consider four cases:

Case 1. $v = u_{1,j}$ $(2 \leq j \leq r)$, that is, u and v are in the same part.

By above coloring, edges $u_{1,1}u_{s,j}$ and $u_{s,j}u_{1,j}$ have distinct colors where $2 \leq s \leq \ell^2$, and so path $u_{1,1}, u_{s,j}, u_{1,j}$ is a rainbow $u - v$ path. Similarly, path $u_{1,1}, u_{s,1}, u_{1,j}$ is a rainbow $u - v$ path for each $2 \leq s \leq \ell^2$. And any two such rainbow paths are disjoint, and so there are at least $2(\ell^2 - 1)$ disjoint rainbow $u - v$ paths in G.

Case 2. $v = u_{s,1}$ $(2 \leq s \leq \ell^2)$, that is, u and v are in the same G_j (here $j = 1$).

Then in subgraph G_1 there are $\ell-1$ disjoint rainbow $u-v$ paths, and in each subgraph G_j where $(2 \leq j \leq r)$, there are $\ell-2$ disjoint rainbow $u_{1,j}-u_{s,j}$ paths of length 2 $u_{1,j}, y, u_{s,j}$ for some ys. Since the color of $u_{1,j}y$, $yu_{s,j}$ is different from that of $u_{1,1}y$, $yu_{s,1}$, respectively, paths $u_{1,1}, y, u_{s,1}$ are $\ell - 2$ disjoint rainbow paths in each G_j where $(2 \leq j \leq r)$. So there are totally $r(\ell-2)+1$ disjoint rainbow $u-v$ paths in G.

Case 3. $v=u_{s,j_0}$ where $2\le s\le \ell, 2\le j_0\le r$, that is, u and v belong to the same $\overline{V_i}$ (here $i=1$) but in distinct parts and subgraphs G_js.

At first, the edge $u_{1,1}u_{s,j_0}$ is a rainbow $u-v$ path.

Next we consider the monochromatic $u_{1,1}-u_{s,1}$ path $u_{1,1}, y, u_{s,1}$. Then each path $u_{1,1}, y, u_{s,j_0}$ is a rainbow $u_{1,1}-u_{s,j_0}$ path of length 2 where $2\le s\le \ell, 2\le j_0\le r$. In U_1, we choose $y=u_{(i-1)\ell+t,1}$ where $1\le i\le \ell, 2\le t\le \ell$ and $t\ne s$. So there are $\ell(\ell-2)$ such paths.

Similarly, there are $\ell(\ell-2)$ disjoint monochromatic $u_{1,j_0}-u_{s,j_0}$ paths u_{1,j_0}, y, u_{s,j_0} where $y\in U_{j_0}$ and we can obtain other $\ell(\ell-2)$ disjoint rainbow $u-v$ paths.

Moreover, in each subgraph G_j where $2\le j\le r$ and $j\ne j_0$, there are $\ell-2$ disjoint rainbow $u_{1,j}-u_{s,j}$ path $u_{1,j}, y, u_{s,j}$ of length 2. Since the color of $u_{1,j}y$ and $yu_{s,j}$ is different from that of $u_{1,1}y$ and yu_{s,j_0}, respectively, paths $u_{1,1}, y, u_{s,j_0}$ are disjoint rainbow $u-v$ paths.

So the number of disjoint rainbow $u-v$ paths in G is $1+2\ell(\ell-2)+(r-2)(\ell-2)=1+(2\ell+r-2)(\ell-2)$.

Case 4. $v=u_{(i_0-1)\ell+t_0,j_0}$ where $2\le i_0\le \ell, 1\le t_0\le \ell, 2\le j_0\le r$, this is, v and u are in distinct $\overline{V_i}$s (here v is not in $\overline{V_1}$).

Subcase 4.1. $t_0=1$.

Then edge uv is a rainbow path. In subgraph G_1, we find the monochromatic $u-u_{(i_0-1)\ell+1,1}$ path $u, y, u_{(i_0-1)\ell+1,1}$. We choose any vertex of $U_1\setminus\{U_{1,1}\cup U_{i,1}\}$ as y. Since the color of edge $yu_{(i_0-1)\ell+1,1}$ is different from that of yv, each path u, y, v is rainbow, and there are $\ell^2-2\ell$ disjoint rainbow $u-v$ paths.

Similarly, in subgraph G_{j_0}, we can find monochromatic paths $u_{1,j_0}, y, u_{(i_0-1)\ell+1,j_0}$, and get other $\ell^2-2\ell$ disjoint rainbow $u-v$ paths.

In each subgraph G_j where $j\ne 1, j_0$, we find the disjoint rainbow $u_{1,j}-u_{(i_0-1)\ell+1,j}$ paths $u_{1,j}, y, u_{(i_0-1)\ell+1,j}$ of length 2. Since the color of $u_{1,j}y$ and $yu_{(i_0-1)\ell+1,j}$ is different from that of $u_{1,1}y$ and $yu_{(i_0-1)\ell+1,1}$, respectively, there are $\ell-2$ disjoint rainbow $u-v$ paths $u_{1,1}, y, u_{(i_0-1)\ell+1,1}$.

So the number of disjoint rainbow $u-v$ paths in G is $1+2\ell(\ell-2)+(r-2)(\ell-2)=1+(2\ell+r-2)(\ell-2)$.

Subcase 4.2. $t_0\ne 1$.

Then, edge uv is a rainbow path. In subgraph G_1, we find the monochromatic $u-u_{(i_0-1)\ell+t_0,1}$ path $u, y, u_{(i_0-1)\ell+t_0,1}$. Let $y=u_{(i-1)\ell+t,1}$ where $i\ne 1, i_0, t\ne 1, t_0$ or $y=u_{t_0,1}$ or $u_{(i_0-1)\ell+1,1}$. Since the color of edge $yu_{(i_0-1)\ell+t_0,1}$ is different from that of yv, each path

u, y, v is rainbow, and so there are $(\ell-2)(\ell-2)+2$ disjoint rainbow $u-v$ paths.

Similarly, in subgraph G_{j_0}, we can find monochromatic paths $u_{1,j_0}, y, u_{(i_0-1)\ell+1,j_0}$, and get other $(\ell-2)(\ell-2)+2$ disjoint rainbow $u-v$ paths.

In each subgraph G_j where $j \neq 1, j_0$, we find the disjoint rainbow $u_{1,j} - u_{(i_0-1)\ell+t_0,j}$ paths $u_{1,j}, y, u_{(i_0-1)\ell+t_0,j}$ of length 2. Since the color of $u_{1,j}y$ and $yu_{(i_0-1)l+t_0,j}$ is different from that of $u_{1,1}y$ and $yu_{(i_0-1)l+t_0,j_0}$, respectively, there are $\ell-2$ disjoint rainbow $u-v$ paths $u_{1,1}, y, u_{(i_0-1)\ell+t_0,j_0}$ (here $y \in G_j$ where $j \neq 1, j_0$).

So the number of disjoint rainbow $u-v$ paths in G is $5+2(\ell-2)(\ell-2)+(r-2)(\ell-2) = 5+(2\ell+r-6)(\ell-2)$.

We know here $\ell \geq 2$, and so $5+(2\ell+r-6)(\ell-2) > r(\ell-2)+1$. Clearly, $1+(2\ell+r-2)(\ell-2) > r(\ell-2)+1$. Let $\ell_0 = \lceil \max\{\sqrt{\frac{k}{2}+1}, \frac{k-1}{r}+2\} \rceil$. We then have $2(\ell_0^2-1) \geq k$, $r(\ell_0-2)+1 \geq k$. So, our following theorem holds.

Theorem 2.2 *For every integer $k \geq 2$, there exists an integer $\ell_0 = \lceil \max\{\sqrt{\frac{k}{2}+1}, \frac{k-1}{r}+2\} \rceil$, such that if $\ell \geq \ell_0$, then $rc_k(K_{\ell^2[r]}) = 2$.* ∎

Next we will consider the case for general complete multipartite graph $G \cong K_{\ell[r]}$ with equal part, where $\ell \geq \ell_0^2$. Let ℓ_1 be the integer satisfying $\ell_1^2 \leq \ell \leq (\ell_1+1)^2$ where $\ell_1 \geq \ell_0$. Then by our Theorem 2.2, we need to consider the case $1 \leq q = \ell - \ell_0^2 \leq 2\ell_1$. Now G can be obtained from $K_{\ell_1^2[r]}$ by adding q new parts P_i $(1 \leq i \leq q)$. Let the vertex set of each corresponding new part be $V(P_i) = \{v_{i,1}, v_{i,2}, \cdots, v_{i,r}\}$ $(1 \leq i \leq q)$. We now update $\overline{V_i}$ as follows: If $1 \leq q \leq \ell_1$, let the new $\overline{V_i}$ be the union of the old $\overline{V_i}$ and $V(P_i)$; If $\ell_1 < q \leq 2\ell_1$, let the new $\overline{V_i}$ be the union of the old $\overline{V_i}$ and $V(P_i) \cup V(P_{\ell_1+i})$ (if there exists) as shown in the Figure 2.1. Similarly, we update U_j $(1 \leq j \leq r)$ (also $G_j, U_{i,j}, G_{i,j}$) as shown in the Figure 2.1 (for example, the new U_1).

Similar to the proof of Theorem 2.1 in [2], we give the 2-edge-coloring of graph G as follows: For each G_j $(1 \leq j \leq r)$, we assign the edge uv of G the color 1 if either $uv \in E(G_{i,j})$ for some i $(1 \leq i \leq \ell_1)$ or if $uv = u_{(i_1-1)\ell_1+t,j}\ u_{(i_2-1)\ell_1+t,j}$ or $uv = v_{i_1,j}\ v_{i_2,j}$ or $uv = v_{\ell_1+i_1,j}\ v_{\ell_1+i_2,j}$ for some i_1, i_2, t with $1 \leq i_1, i_2, t \leq \ell_1$ and $i_1 \neq i_2$. All other edges of G_j are assigned the color 2.

For other edges, that is, edges between distinct G_js, we use above two colors as follows: If $uv = u_{i_1,j_1}\ u_{i_2,j_2}$ where $u_{i_1,j_1} \in G_{j_1}$, $u_{i_2,j_2} \in G_{j_2}$, we give it the color different from that of edge $u_{i_1,j_1}\ u_{i_2,j_1}$ of graph G_{j_1} or $u_{i_1,j_2}\ u_{i_2,j_2}$ of graph G_{j_2} (since edges $u_{i_1,j_1}\ u_{i_2,j_1}$ and $u_{i_1,j_2}\ u_{i_2,j_2}$ have the same color by above coloring); Similarly, color the edge for the cases that $uv = u_{i_1,j_1}v_{i_2,j_2}$ and $uv = v_{i_1,j_1}v_{i_2,j_2}$.

By the proof of our Theorem 2.2, there are at least k disjoint rainbow paths connecting any two vertices $u, v \in V(G) \setminus \bigcup_{i=1}^{q} V(P_i)$. We need to consider the paths between u, v



Figure 2.1 The figure of Theorem 2.4.

where $u \in \bigcup_{i=1}^{q} V(P_i)$ and $v \in V(G)$. Without loss of generality, let $u = v_{1,1}$, and we consider four cases, which are similar to the four cases in the proof of our Theorem 2.2.

Case 1. $v \in V(P_1)$, that is, u and $v = v_{1,j_0}$ $2 \leq j_0 \leq r$ are in the same part (here the part is P_1).

G contains the disjoint rainbow $u-v$ paths $v_{1,1}, y, v_{1,j_0}$, and we let y be the elements of $U_1 \cup U_{j_0} \setminus \{v_{1,1}, v_{1,j_0}\}$. So the number of these paths is at least $2\ell_1^2 \geq 2\ell_0^2 > 2(\ell_0^2 - 1) \geq k$.

Case 2. $v \in U_1$, that is, v and u are in the same subgraph G_1,

In G_1, there are at least $\ell_1 - 1$ rainbow $u - v$ paths; In each G_j $(2 \leq j \leq r)$ there are at least $\ell_1 - 2$ rainbow $u' - v'$ paths of length 2: u', y, v' where u', v' is in the same part as u, v, respectively, and we get at least $\ell_1 - 2$ rainbow $u - v$ paths: u, y, v, where $y \in G_j$ $(2 \leq j \leq r)$.

So the number of disjoint rainbow $u-v$ paths is at least $1+r(\ell_1-2) \geq 1+r(\ell_0-2) \geq k$.

Case 3. $v \in \overline{V_1}$ but not in $U_1 \cup V(P_1)$. Without loss of generality, let $v \in U_{j_0}$.

Then, edge uv is a $u - v$ path.

Subcase 3.1. $v = v_{l+1,j_0}$.

Then similar to Case 3 in the proof of of Theorem 2.2, we find the monochromatic $u - v_{l+1,1}$ path of length 2 in subgraph G_1 and get at least ℓ_1^2 disjoint rainbow $u - v$ paths. Similarly, there are other ℓ_1^2 disjoint rainbow $u - v$ paths by finding the monochromatic $v_{1,j_0} - v_{\ell+1,j_0}$ paths in G_{j_0}.

In each G_j $(j \neq 1, j_0)$, similar to Case 3 in the proof of Theorem 2.2, we get at least $\ell_1 - 2$ disjoint rainbow $u - v$ paths of length 2 by finding rainbow $v_{1,j} - v_{\ell+1,j}$ paths in G_j.

So the number of disjoint rainbow $u - v$ paths is at least $1 + 2\ell_1^2 + (r-2)(\ell_1 - 2) > 1 + 2\ell_1(\ell_1 - 2) + (r-2)(\ell_1 - 2) = 1 + (2\ell_1 + r - 2)(\ell_1 - 2) \geq 1 + (2\ell_0 + r - 2)(\ell_0 - 2) \geq k$.

Subcase 3.2. $v = u_{s_0,j_0}$ where $1 \leq s_0 \leq \ell_1, 2 \leq j_0 \leq r$.

With a similar procedure to Subcase 3.1, we can get at least $1 + 2\ell_1(\ell_1 - 1) + (r - 2)(\ell_1 - 2) > 1 + (2\ell_1 + r - 2)(\ell_1 - 2) \geq 1 + (2\ell_0 + r - 2)(\ell_0 - 2) \geq k$ rainbow disjoint $u - v$ paths in graph G.

Case 4. v is not in $\overline{V_1}$ and U_1, without loss of generality, let $v \in \overline{V_2} \setminus U_1$. We consider two subcases similar to Case 4 in the proof of Theorem 2.2.

Case 4.1. $v = v_{2,j_0}$ where $2 \leq j_0 \leq r$.

Similar to Subcase 4.1 in the proof of Theorem 2.2, we can get at least $1 + 2\ell_1(\ell_1 - 2) + (r-2)(\ell_1 - 2) = 1 + (2\ell_1 + r - 2)(\ell_1 - 2) \geq 1 + (2\ell_0 + r - 2)(\ell_0 - 2) \geq k$ disjoint rainbow $u - v$ paths in G.

Case 4.2. $v \neq v_{2,j_0}$.

Similar to Subcase 4.2 in the proof of Theorem 2.2, we can get at least $1 + 2(\ell_1 - 1)(\ell_1 - 2) + (r - 2)(\ell_1 - 2) = 1 + (2\ell_1 + r - 4)(\ell_1 - 2) \geq 1 + r(\ell_1 - 2) \geq 1 + r(\ell_0 - 2) \geq k$ (since $\ell_1 \geq 2$) disjoint rainbow $u - v$ paths in G.

With above discussion, we have our following theorem:

Theorem 2.3 *For every integer $k \geq 2$, there exists an integer $\ell_0 = \lceil \max\{\sqrt{\frac{k}{2} + 1}, \frac{k-1}{r} + 2\}\rceil$, such that if $\ell \geq \ell_0^2$, then $rc_k(K_{\ell[r]}) = 2$.* ∎

We now obtain a new graph G' from G by adding p new vertices: $w_1, w_2, \cdots, w_p$ $(1 \leq p \leq r - 1)$ and edge vw_j where $1 \leq j \leq p$ and $v \in V(G)$. Color the new edges as follows: Give color 2 to edges between U_j and w_j, and color 1 to other edges.

Now we count the number of rainbow disjoint $u - v$ paths between any two vertices of G. We need to consider the case that $u = w_j, v \in V(G')$ $(1 \leq j \leq p)$. Without loss of generality, let $u = w_1$.

Case 1. $v \in V(G)$.

Subcase 1.1. $v \in U_1$, without loss of generality, let $v = u_{1,1}$.

Then G contains the $u - v$ path u, v as well as the $u - v$ rainbow paths $w_1, y, u_{1,1}$ where $y \in \overline{V_1} \setminus \{u_{1,1}, \cdots, u_{1,r}\}$, or $y = u_{(i-1)\ell_1+1,j}(2 \leq i \leq \ell_1, 1 \leq j \leq r)$. So the number of disjoint rainbow $u - v$ paths is at least $1 + 2r(\ell_1 - 1) > 1 + r(\ell_0 - 2) \geq k$.

Subcase 1.2. v is not in U_1, without loss of generality, let $v = u_{1,2}$.

Then G contains the $u - v$ path u, v as well as the $u - v$ rainbow paths $w_1, y, u_{1,2}$ where $y \in (U_1 \cup U_2) \setminus (U_{1,1} \cup U_{1,2} \cup \{u_{(i-1)\ell_1+1,j_0}\}_{i=1}^{\ell_1})$ where $j_0 = 1, 2$, or $y \in \{u_{(i-1)i_1+1,j_1}\}_{i=2}^{\ell_1}$ where $j_1 \neq 1, 2$, or $y = u_{s,j}$ where $2 \leq s \leq \ell_1, j \neq 1, 2$. So the number of disjoint rainbow $u - v$ paths is at least $1 + 2(\ell_1 - 1)(\ell_1 - 1) + 2(\ell_1 - 1)(r - 2) = 1 + (\ell_1 - 1)(2\ell_1 + 2r - 6) > 1 + (\ell_1 - 2)(2\ell_1 + 2r - 6) \geq 1 + (\ell_1 - 2)r \geq k$(here we let $r \geq 2$).

Case 2. $v = w_j$ where $j \neq 1$.

Then G contains the $u - v$ rainbow paths w_1, y, w_j where $y \in U_1 \cup U_j$. So the number of disjoint rainbow $u - v$ paths is at least $2\ell_1^2 \geq 2\ell_0^2 > 2(\ell_0^2 - 1) \geq k$.

If $r = 1$, then $G' = G$ is complete, so by Theorem 2.1 of [2] and above discussion, we have the following theorem:

Theorem 2.4 *Let G be a complete $(\ell + 1)$-partite graph with ℓ parts of size $r \geq 1$ and a part of size p where $1 \leq p < r$. Then for every integer $k \geq 2$, if $\ell \geq \ell_0^2$ where $\ell_0 = \lceil \max\{\sqrt{\frac{k}{2} + 1}, \frac{k-1}{r} + 2\}\rceil$, then $rc_k(G) = 2$.* ∎

We know that the graph in Theorem 2.4 is a spanning subgraph of the complete K_n with $n = \ell r + p$ and $\ell \geq \ell_0^2$. Now we let $r = r_0 = \lceil \sqrt{2k} \rceil$, and G be the complete graph K_n with $n \geq \ell_0^2 r_0$. Then, since $n = \ell_3 r_0 + p$ where $\ell_3 \geq \ell_0^2, 0 \leq p < r_0$, by Theorem 2.4 we have

$rc_k(G)=2$, here $\ell_0=\lceil\max\{\sqrt{\frac{k}{2}+1},\frac{k-1}{r}+2\}\rceil=\lceil\max\{\sqrt{\frac{k}{2}+1},\frac{k-1}{\lceil\sqrt{2k}\rceil}+2\}\rceil=\lceil\frac{k-1}{\lceil\sqrt{2k}\rceil}\rceil+2$. Thus we have

Corollary 2.5 *For every integer $k\geq 2$, let integer $f(k)=\ell_0^2 r_0$. If $n\geq f(k)$, then $rc_k(K_n)=2$, where $\ell_0=\lceil\frac{k-1}{\lceil\sqrt{2k}\rceil}\rceil+2$ and $r_0=\lceil\sqrt{2k}\rceil$.* ∎

From Corollary 2.5, we know that for the integer $f(k)$, we have that $f(k)\leq ck^{\frac{3}{2}}+C$ where $0<c<1$ is a constant and $C=o(k^{\frac{3}{2}})$, and this upper bound is a considerable improvement of the bound given in Theorem 2.1 of [2].

References

[1] J.A. Bondy, U.S.R. Murty, *Graph Theory*, GTM 244, Springer, 2008.

[2] G. Chartrand, G.L. Johns, K.A. McKeon, P. Zhang, The rainbow connectivity of a graph, accepted for publication in *Networks – An International Journal.*

[3] A.B. Ericksen, A matter of security, *Graduating Engineer & Computer Careers* (2007), 24-28.